

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/269847156>

A COMPARATIVE STUDY OF MALONDIALDEHYDE CONTENTS OF SOME MEAT AND FISH SAMPLES PROCESSED BY DIFFERENT METHODS

Article · July 2013

DOI: 10.7897/2277-4572.02448

CITATIONS

3

READS

74

2 authors, including:



Ngozi Paulinus Okolie
University of Benin

60 PUBLICATIONS 515 CITATIONS

[SEE PROFILE](#)

Some of the authors of this publication are also working on these related projects:



Multifunctional nanosystem based on natural proteins for targeted delivery of anti-cancer agent to Colorectal Cancer cells [View project](#)



Toxicity studies [View project](#)



A COMPARATIVE STUDY OF MALONDIALDEHYDE CONTENTS OF SOME MEAT AND FISH SAMPLES PROCESSED BY DIFFERENT METHODS

Okolie Ngozi Paulinus^{1*} and Okugbo, Osarhieme Tinuade²

¹Department of Biochemistry, Faculty of Life Sciences, University of Benin

²Department of Biochemistry, Benson Idahosa University, Benin City, Nigeria

*Corresponding Author Email: Okolie_ngozi@yahoo.co.uk

DOI: 10.7897/2277-4572.02448

Published by Moksha Publishing House. Website www.mokshaph.com

All rights reserved.

Received on: 24/06/13 Revised on: 25/07/13 Accepted on: 29/07/13

ABSTRACT

The effect of different processing methods on the malondialdehyde, MDA contents of 7 popular samples of fresh meat and fish eaten in Nigeria, was investigated. The MDA levels were assayed in a colorimetric reaction with thiobarbituric acid in fresh samples of beef, pork, turkey, chicken, snail, catfish and goat meat processed by boiling, frying, roasting and freezing. Results obtained showed that all processing methods led to significant increases in MDA contents of the samples studied relative to their fresh, unprocessed counterparts ($p < 0.05$). For most samples, boiling, roasting and freezing yielded comparable levels of MDA. However, significantly higher levels of MDA were seen in samples fried after boiling, relative to boiled samples ($p < 0.05$). For the boiled samples, beef and goat meat had the least MDA. The order of MDA content in the boiled samples was beef > goat meat > catfish > turkey > chicken > snail > pork. The MDA content of frozen mackerel was significantly increased by boiling and frying and by further freezing ($p < 0.05$). These results suggest that while all the processing methods may be unsafe, the frying of boiled meat and fish may be particularly deleterious to consumer health, in view of the well-known mutagenic and carcinogenic effects of MDA.

Keywords: Malondialdehyde Content, Meat, Fish, Snail, Boiling, Roasting, Frying.

INTRODUCTION

Lipid peroxidation is regarded as a major source of quality deterioration, not only in meat and meat products but also in fish and fish products. Aside from the producing undesirable changes in flavor and loss of food quality, lipid peroxidation increases the level of malondialdehyde, MDA in foods. Studies have revealed that MDA is one of the most abundant lipid peroxidation cytotoxins formed in foods, especially in meat^{1,2}. Earlier reports on the mutagenic and carcinogenic potentials of MDA^{3,4} have since led to increased interest in its implication for human health. Following ingestion of peroxidised foods, humans and lower animals have been shown to excrete increased amounts of MDA in the urine⁵. Indeed it is now known that MDA can be absorbed from tainted foods when these foods are injected^{6,7}. MDA is genotoxic, reacting with DNA to form highly mutagenic adducts in human cells⁸⁻¹⁰. It has been demonstrated that the level of MDA-DNA adducts in human colorectal tissue correlates with diet and incidence of adenoma¹¹. Thus the presence of MDA in foods may have deleterious consequences for human health. In Nigeria as is the case in most other countries of the world, meat and fish are the major sources of dietary protein. Both are usually consumed after thermal processing which may entail boiling, frying or roasting/grilling. From toxicological perspectives, fresh meat and fish are safer than frozen ones, because it has been demonstrated that freezing results in increased accumulation of MDA in meat and fish¹². Thus thermal treatment of fresh samples may be safer. However prolonged heating is known to also increase the level of lipid peroxidation by a combination of several processes including disruption of muscle cell structure, inactivation of anti-oxidant enzymes and release of oxygen and iron from myoglobin¹⁵. The present study attempts to compare the effects of different thermal processing methods on the MDA contents of popular meat and fish samples consumed in Nigeria, with a view to

determining the safest procedure, in view of the toxic implications of MDA.

MATERIALS AND METHODS

Sample Collection and Processing

Lean muscle samples of beef, pork and goat meat were purchased from freshly sacrificed cow, pig and goats respectively, in an abattoir in Benin City, Edo State, Nigeria. Live African giant snail, live chicken and live turkey were procured from Oba Market, Benin City, while live catfish was obtained from Benson Idahosa University Farms, Benin City. Frozen fish (mackerel) was purchased from a retail outlet in a cold room at Oba market, Benin City. The live samples were sacrificed and muscle parts were dissected out. Muscle was also collected from frozen mackerel. All samples were washed clean with tap water and divided into two portions. One portion was used immediately for analysis of MDA contents, while MDA was analysed in the other portion after processing by boiling, frying + roasting, and freezing. The samples were sliced into small sizes similar to those routinely used in cooking, roasting and frying. For processing by boiling, the samples were immersed in boiling water in clean aluminum pots and allowed to cook for exactly 15 minutes before removal. A part of each boiled sample was further processed by deep-frying for 5 minutes in Gino^R brand vegetable oil. For processing by roasting, the sliced fresh samples were wrapped in clean aluminum foil and divided into two parts. One part was roasted for 1 h at 170^o C in an oven, while the other part was stored in a deep freezer at -10^o C for 5 days. All processed samples were subsequently analysed for their MDA contents.

Analysis of Samples for MDA

Fresh and processed samples were analysed for MDA using the colorimetric reaction with thiobarbituric acid, TBA¹⁴. In essence, 0.5 g of sample was thoroughly ground with some

grains of acid-washed sand in a hand mortar and extracted with 10 ml of 15 % TCA for 15 minutes. The resultant homogenate was centrifuged at 3000 rpm for 10 minutes and the supernatant fraction subjected to MDA assay. The reaction mixture contained 1 ml of extract and 2 ml of TBA reagent. The mixture was heated for 15 minutes in a boiling water bath, and on cooling; the resultant precipitate was removed by centrifugation at 3000 g for 10 minutes. Then the absorbance of the pink-colored solution was read at 535 nm and converted to moles of MDA by using its molar extinction coefficient. MDA levels were expressed as mole/g fresh wt.

Statistics

The levels of MDA (Mean ± SD) arising from the various processing methods were each compared with corresponding

values for unprocessed samples using student’s t-test. P values < 0.05 were taken as significant. T-test was also used to compare values obtained for boiled samples and boiled + fried samples; while analysis of variance, ANOVA was used to compare differences between frozen mackerel MDA values arising from different processing methods.

RESULTS

Table 1 shows the levels of MDA in fresh and processed samples of beef, pork, catfish, snail, chicken, turkey and goat meat. MDA was lowest in the fresh samples, but increased with boiling and frying, except in snail where boiling actually led to a decrease in MDA. For most samples, boiling and roasting produced comparable levels of MDA. The highest increases in MDA were seen in boiled and fried samples.

Table 1: MDA Contents of Fresh and Processed Samples of Catfish, Goat meat, Beef, Chicken, Turkey, Pork and Snail

Sample / Processing method	MDA Content (mole/g fresh wt x 10 ⁻³)				
	Fresh	Boiled	Boiled and deep-fried	Roasted	Frozen
Catfish	1.71 ± 0.20 ^a	2.30 ± 0.26 ^b	3.33 ± 0.46 ^c	2.91 ± 0.20 ^c	2.65 ± 0.2 ^c
Beef	0.13 ± 0.04 ^a	1.62 ± 0.20 ^b	4.19 ± 0.39 ^c	2.26 ± 0.27 ^d	2.44 ± 0.13 ^d
Snail	4.74 ± 1.11 ^a	3.72 ± 0.97 ^b	8.99 ± 1.32 ^c	4.55 ± 0.64 ^d	5.81 ± 0.65 ^d
Turkey	0.38 ± 0.20 ^a	2.38 ± 0.35 ^b	5.00 ± 0.46 ^c	3.02 ± 0.42 ^d	2.69 ± 0.34 ^b
Chicken	0.60 ± 0.15 ^a	2.46 ± 1.02 ^b	6.48 ± 0.35 ^c	3.12 ± 0.20 ^d	3.95 ± 0.34 ^c
Pork	1.41 ± 0.13 ^a	3.80 ± 0.58 ^b	7.31 ± 0.68 ^c	4.74 ± 0.68 ^d	2.91 ± 0.49 ^c
Goat meat	0.26 ± 0.07 ^a	1.76 ± 0.26 ^b	4.49 ± 0.13 ^c	2.31 ± 0.13 ^d	1.92 ± 0.26 ^b

Results are expressed as Mean ± SD (n = 3). For each sample, values across bearing superscripts different from the corresponding value for fresh sample differ significantly from it (p < 0.5).

Samples subjected to dual heat treatment of boiling and frying had at least twice as much % increase in MDA over fresh samples as boiled samples (Table 2). This trend was

also seen in snail, where boiling and frying brought about significant increases in MDA levels in spite of the fact that boiling alone actually decreased its MDA content (Table 1).

Table 2: Increase in Mean MDA Levels of Boiled Samples and Boiled + Fried Samples Relative to Unprocessed Samples (%)

Sample	Boiled	Boiled + fried
Catfish	34	130
Beef	1146	3123
Snail	-22	89
Turkey	526	1216
Chicken	310	980
Pork	170	418
Goat meat	577	1627

Boiling significantly reduced the MDA content of raw, frozen mackerel, but with boiling and frying, the MDA level was significantly increased relative to the value for the raw

sample (p < 0.05) (Table 3). Roasting and further freezing also led to significant increases in the MDA content of the raw, frozen mackerel (p < 0.05).

Table 3: Effect of Various Processing Methods on the MDA Content of Frozen Mackerel

Frozen mackerel	MDA (mole/g fresh wt x 10 ⁻³)
Raw	9.61 ± 0.38 ^a
Boiled	6.79 ± 0.38 ^b
Boiled and fried	11.97 ± 0.45 ^c
Roasted	11.80 ± 0.71 ^c
Further freezing	10.81 ± 0.51 ^c

Values are Mean ± SD; n=3. Values with different superscripts differ significantly (p < 0.05)

DISCUSSION

The results obtained in this study show that heat and cold treatments accentuate levels of MDA in fish and meat. It has been reported that heating increases the level of lipid peroxidation in meat by a number of processes including inactivation of protective antioxidant enzymes, disruption of muscle cell structure and release of oxygen and iron from myoglobin¹⁴. Iron has been implicated as the most probable catalyst in the promotion of lipid peroxidation through generation of hydroxyl radicals by the Fenton reaction^{15,16}.

Most muscle foods contain myoglobin and hemoglobin-bound iron, which contribute 74, 47 and 28 % of total iron in beef, pork and chicken thigh respectively¹⁷. In addition, the oxidative stability of meat is dependent on the degree of unsaturation of its sub-cellular membrane phospholipids: a high degree of unsaturation is associated with higher oxidative potentials¹⁸. Although snail is low in fat, about 75 % of the fat is composed of polyunsaturated fatty acids, which render it vulnerable to lipid peroxidation¹⁹. Thus the differences in levels of MDA among the samples studied may

be a reflection of the differences in their susceptibilities to lipid peroxidation. In general it has been demonstrated that the extent of lipid peroxidation in meat depends on the anatomical location of the muscle, muscle type and animal species^{20,21}. Freezing significantly increased the MDA contents of all the fresh samples of meat and catfish relative to fresh samples and in most cases relative to boiled samples also. This is in agreement with previous reports that freezing increases MDA contents of fish and meat^{12,22,23}. Although the precise mechanism by which refrigeration increases lipid peroxidation is not well understood, it may not be unconnected with disruption of membrane structure and exposure of phospholipids to pro-oxidants through slicing as well as freezing and thawing. Fatty fish such as mackerel have high oxidation potentials, even when stored at sub-zero temperatures²³. This explains the high level of MDA in frozen mackerel and the fact that the MDA increased significantly with further freezing. The levels of MDA resulting from roasting (grilling) were in most samples comparable with MDA levels from boiling. This is surprising, considering the fact that the samples were grilled for 1 hour while boiling lasted exactly 15 minutes. It is possible that the higher grilling temperature of 150°C resulted in lower rate of lipid peroxidation due to rapid dehydration of the samples, unlike boiling which took place in a hydrated medium and at a lower temperature. Boiling resulted in the lowest levels of MDA in all samples and thus appears to be the safest of all the processing methods investigated. It also resulted in significant reduction of MDA in frozen mackerel and was the only processing method that was able to achieve this. The reduction of MDA content of frozen mackerel by boiling is considered a crucial finding, because most Nigerian homes and hotels depend on frozen meat and fish, which contain appreciable levels of MDA¹². It is interesting that samples fried after boiling had significantly higher level of MDA when compared to MDA resulting from any other treatment. This may be explained by the fact that the samples were exposed to two different heat treatments, each of which is known to enhance lipid peroxidation. Frying of boiled meat and fish are rather popular culinary habits in most Nigerian homes and public eating places. From the results obtained in this study, it appears that this practice may be deleterious to consumer health due to rapid build-up of MDA. The presence of MDA in foods is of considerable health concern to the consumer, in view of its well-known carcinogenic and mutagenic effects²⁴. MDA is classified as perhaps the most abundant cytotoxin arising from lipid peroxidation in muscle-derived foods, especially meat^{1,2}. MDA is readily absorbed from ingested food in the form of N-(2-propenal)-lysine⁶. Indeed N-2-propenal lysine derivative is the major form of MDA in the urine following consumption of peroxidised foods². In this form, MDA still retains its ability to modify DNA. Thus absorbed MDA forms adducts with deoxyadenosine and deoxyguanosine, leading to DNA damage which may pose serious health problems to the consumers²⁵. Consequently processing methods that lead to reduction of MDA in muscle foods are highly desirable. Since meat and fish are rarely eaten raw, our results show that boiling of these vital sources of dietary protein may be the safest method of cooking them in homes and public places.

CONCLUSION


The results of this study have shown that the practice of frying meat and fish after boiling, as well as the popular habit of protracted refrigeration should be discouraged, in view of the significant elevation of lipid peroxidation associated with them. Boiling appears to be the safest way to process muscle foods.

REFERENCES

1. Stocker R, Kearney JF Jr. New insights on oxidative stress in the artery wall. *J Thromb Haemost* 2005; 3: 1825-1834. <http://dx.doi.org/10.1111/j.1538-7836.2005.01370.x> PMID:16102049
2. Gorelik S, Kohan R, Ligumsky M, Kanner J. Saliva plays a dual role in oxidation process in stomach medium. *Arch Biochem Biophys* 2007; 458: 236-243. <http://dx.doi.org/10.1016/j.abb.2006.12.006> PMID:17250799
3. Shamberger R, Andreone T, Willis C. Antioxidants and cancer. IV. Initiating activity of malondialdehyde as a carcinogen. *J Natl Cancer Inst* 1974; 53: 1771-1773. PMID:4436871
4. Mukai FH, Goldstein BD. Mutagenicity of malondialdehyde, a decomposition product of peroxidised polyunsaturated fatty acids. *Science* 1976; 191: 868-869. <http://dx.doi.org/10.1126/science.766187> PMID:766187
5. Hadley M, Draper HH. Identification of N-(2-propenal) ethanolamine as a urinary metabolite of malondialdehyde. *Free Rad Biol Med* 1989; 6: 49-52. [http://dx.doi.org/10.1016/0891-5849\(89\)90159-7](http://dx.doi.org/10.1016/0891-5849(89)90159-7)
6. Gopaul NK, Halliwell B, Anggard EE. Measurement of plasma F2 isoprostanes as an index of lipid peroxidation does not appear to be compounded by diet. *Free Rad Res* 2000; 33: 115-127. <http://dx.doi.org/10.1080/1071576000300671> PMID:10885619
7. Giron Calle J, Alaiz M, Millan M, Ruiz Gutierrez V, Vioque E. Bound malondialdehyde in foods: bioavailability of the N-2-propenals of lysine. *J Agric Food Chem* 2002; 50: 6194-6198. <http://dx.doi.org/10.1021/jf025681r> PMID:12358501
8. Riggins JN, Marnett LJ. Mutagenicity of the malondialdehyde oligomerization products 2-(3-oxo-1-propenyl) malondialdehyde and 2,4-dihydroxymethylene -3-(2,2-dimethoxyethyl) glutaraldehyde in *Salmonella*. *Mut Res* 2001; 497: 153-157. [http://dx.doi.org/10.1016/S1383-5718\(01\)00253-4](http://dx.doi.org/10.1016/S1383-5718(01)00253-4)
9. Cline SD, Riggins JN, Tornaletti S, Marnett LJ, Hanawalt PC. Malondialdehyde adducts in DNA arrest transcription by T7 RNA polymerase and mammalian RNA polymerase II. *Proc Natl Acad Sci* 2004; 101: 7275-7280. <http://dx.doi.org/10.1073/pnas.0402252101> PMID:15123825 PMID:PMC409909
10. Del Rio D, Stewart AJ, Pellegrini N. A review of recent studies on malondialdehyde as toxic molecule and biological marker of oxidative stress. *Nutr Metab Cardiovasc Dis* 2005; 15: 316-328. <http://dx.doi.org/10.1016/j.numecd.2005.05.003> PMID:16054557
11. Leuratti C, Watson MA, Deag EJ, Welch A, Singh R, Gottschala E, Marnett LJ, Atkin W et al. Detection of malondialdehyde DNA adducts in human colorectal mucosa: Relationship with diet and the presence of adenomas. *Cancer Epidemiol Biomarkers Prev* 2002; 11: 267-273. PMID:11895876
12. Okolie NP, Akioyamen MO, Okpoba N, Okonkwo C. Malondialdehyde levels of frozen fish, chicken and turkey on sale in Benin City markets. *Afr J Biotechnol* 2009; 8: 6638-6640.
13. Min E, Ahn DU. Mechanism of lipid peroxidation in meat and meat products – A review. *Food Sci Biotechnol* 2005; 14: 152-163.
14. Buege J, Aust SD. *Microsomal Lipid Peroxidation*. In: Colowick SP, Kaplan NO (eds). *Methods in Enzymology*: Academic Press, New York; 1978. p. 302-311 PMID:672633
15. O Connell M, Halliwell B, Moorehouse CP, Aruoma OI, Baum H, Peter TJ. Formation of hydroxyl radicals in the presence of ferritin and haemosiderin: Is haemosiderin formation a biological protective mechanism? *Biochem J* 1986; 234: 727-731. PMID:3718494 PMID:PMC1146631
16. Liochev SI, Fridovich I. The role of O₂ in the production of OH⁻ *in vitro* and *in vivo*. *Free Rad Biol Med* 1994; 16: 97-112. [http://dx.doi.org/10.1016/0891-5849\(94\)90239-9](http://dx.doi.org/10.1016/0891-5849(94)90239-9)
17. Shriker BR, Miller DD. Effects of cooking and chemical treatment on heme and non-heme iron in meat. *J Food Sci* 1983; 48: 1340-1349. <http://dx.doi.org/10.1111/j.1365-2621.1983.tb09225.x>
18. Coetzee GJM, Hoffman LC. Effect of dietary vitamin E on the performance of broiler meat during refrigerated and frozen storage. *South Afr J Anim Sci* 2001; 31: 158-172. <http://dx.doi.org/10.4314/sajas.v31i3.3799>
19. Su XQ, Antonas KN, Li D. Comparison of N-3 polyunsaturated fatty acid contents of wild and cultured Australian abalone. *Int J Food Sci*

- Nutr 2004; 55: 149-154. <http://dx.doi.org/10.1080/09637480410001666469> PMID:14985187
20. Rhee KS, Ziprin, YS. Lipid peroxidation in retail beef, pork and chicken muscles as affected by concentrations of heme pigments and non-heme iron and microsomal enzymic lipid peroxidation activity. *J Food Biochem* 2002; 11: 1-15. <http://dx.doi.org/10.1111/j.1745-4514.1987.tb00109.x>
21. Rhee KS, Anderson LM, Sams AR. Lipid peroxidation potential of beef, chicken and pork. *J Food Sci* 1996; 61: 8-12. <http://dx.doi.org/10.1111/j.1365-2621.1996.tb14714.x>
22. Eun JB, Boyle JA, Hearnberger JO. Lipid peroxidation and chemical changes in catfish (*Ictalurus punctatus*) muscle microsomes during frozen storage. *J Food Sci* 1994; 59: 251-255. <http://dx.doi.org/10.1111/j.1365-2621.1994.tb06941.x>
23. Olafsdottir G, Martinsdottir E, Oehlenschlager J, Dalgaard P, Jensen B, Undeland I. Methods to evaluate fish freshness in research and industry. *Trends Food Sci Technol* 1997; 8: 258-265. [http://dx.doi.org/10.1016/S0924-2244\(97\)01049-2](http://dx.doi.org/10.1016/S0924-2244(97)01049-2)
24. Niedernhofer LJ, Daniels JS, Rouzer CA, Greene RE, Marnett LJ. Malondialdehyde, a product of lipid peroxidation, is mutagenic in human cells. *J Biol Chem* 2003; 278: 426-433. <http://dx.doi.org/10.1074/jbc.M212549200> PMID:12775726
25. West JD, Marnett LJ. Alterations in gene expression induced by the lipid peroxidation product, 4-hydroxy-2-nonenal. *Chem Res Toxicol* 2005; 18: 1642-1653. <http://dx.doi.org/10.1021/tx050211n> PMID:16300372

Source of support: Nil, Conflict of interest: None Declared



How to cite this article:

Okolie Ngozi Paulinus and Okugbo, Osarhieme Tinuade. A comparative study of Malondialdehyde contents of some meat and fish samples processed by different methods. *J Pharm Sci Innov.* 2013; 2(4): 26-29.